

ULTRA HIGH ENERGY COSMIC RAYS

PASQUALE BLASI

INAF/Osservatorio Astrofisico di Arcetri, Largo E. Fermi, 5 Firenze (Italy)
E-mail: blasi@arcetri.astro.it

The detection of cosmic rays with energy around and in excess of 10^{20} eV raises many questions that future experiments will help answering to. I address here my view of some of these open issues, as they are now and as they might be affected by future observations.

1. Introduction

In the eighty years of history of cosmic rays, there has been a constant search for the end of the cosmic ray spectrum. It has long been thought that this end of the spectrum would be determined by the highest energy that cosmic accelerators might be able to achieve. Despite this continuous search, no end of the spectrum was found. In 1966, right after the discovery of the cosmic microwave background (CMB), it was understood ¹ that high energy protons would inelastically scatter the photons of the CMB and produce pions. For homogeneously distributed sources this would cause a flux suppression, called the *GZK cutoff*: for the first time the end of the cosmic ray spectrum was related to a physical process rather than to speculations on the nature of the accelerators. Moreover, for the first time, the end of the cosmic ray spectrum was predicted to be at a rather well defined energy, around 10^{20} eV, where the so-called photopion production starts to be kinematically allowed. Forty years later, we are still seeking a confirmation that the cosmic ray spectrum has in fact such a flux suppression. The two largest experiments currently operating in the energy range of interest, namely AGASA ² and HiRes ³, appear to have discrepant results in the highest energy end of the spectrum. While the data collected by the former appear to be consistent with the extension of the lower energy spectrum, the latter experiment suggests that the GZK feature may be present in the data. The small statistics of events however does not allow to draw a definitive conclusion about the detection of the GZK feature ⁴. A positive detection would be a proof of the extragalactic origin of UHECRs.

The small statistics of events also affects our ability to assess the significance of another piece of information, that comes from the AGASA experiment: some of the events are clustered on angular scales comparable with the angular resolution of AGASA ⁵. If confirmed, this would represent the first true indication that UHECRs are in fact accelerated in some still unknown powerful astrophysical objects.

Much progress has been made in the understanding of the physics behind the acceleration processes that may be able to energize cosmic rays to the highest observed energies. This is particularly true for shock acceleration, that still remains the most promising candidate acceleration process, probably because it is the one that has been studied the most.

In this short review, after summarizing the main observational facts (Section 2), I will discuss the main open questions that seek an answer (Section 3).

2. A short summary of the observations

In this section I briefly summarize the main pieces of the puzzle of UHECRs:

Isotropy: The directions of arrival of the events at energies above $\sim 4 \times 10^{19}$ eV appear isotropically distributed in the sky. No immediate association with local structures (galactic disc, supergalactic plane) arises from the data.

Lack of source identification: No association of the observed events with known powerful nearby sources has been found. This may represent a serious problem for the highest energy events, with energy higher than 10^{20} eV, for which the loss length is small and the sources are forced to be closeby. However, even at 4×10^{19} eV the loss length becomes of the same order of magnitude of the size of the universe and it is therefore difficult to find a counterpart, in particular because of the poor angular resolution of current experiments.

Small Scale Anisotropies (SSA): The AGASA data show several doublets and triplets of events on angular scales comparable with the resolution of the instrument ⁵. The statistical significance of these multiplets is still the subject of some debate [see for instance ⁶], but if confirmed as not just the result of statistical fluctuations they could in fact represent the first evidence that UHECRs are accelerated in astrophysical point sources. This evidence would point against most so-called top-down models, in which the emission is truly diffuse.

The composition: At the highest energies the information about the chemical composition is so far very poor. A reanalysis of the Haverah Park inclined showers allowed to constrain the fraction of gamma rays at energy larger than 4×10^{19} eV to about 50% ⁷ (see the contribution of Alan Watson in these proceedings ⁸).

3. Open questions

In this section I address some questions that seem to me as particularly important for the understanding of the origin of UHECRs. The list is not supposed to be complete and should only be considered as a possible starting point for further investigations.

3.1. Should we expect a sharp GZK cutoff?

For a long time, the issue of detecting the GZK flux suppression has been interpreted as the search for cosmic ray events with energy above 10^{20} eV and of nearby sources

of these particles. It is often forgotten that the GZK feature is not avoided at all if the source density is constant (homogeneous distribution of the sources) all over the universe. As already mentioned above, a flux suppression is expected due to the fact that above the threshold for photopion production protons lose energy rapidly. It is worth stressing however that what has been named the GZK cutoff is in fact




Figure 1. Loss length of UHECRs for proton pair production and photopion production.

a *feature* that can be more or less pronounced depending upon details such as the injection spectrum of cosmic rays, the luminosity evolution of the sources, the local overdensity of sources and the magnetic field strength in the intergalactic medium. In general terms, the GZK feature is not determined by the number of events above 10^{20} eV but rather by the relative fraction of events above 10^{20} eV compared with that at lower energies. This is easily explained through Fig. 1, where I plot the loss length for proton pair production and photopion production (solid curve) as a function of energy⁹. The straight horizontal line represents the size of the universe today, while the dashed lines are just there to drive the eye to identify the energies of 10^{20} eV and 5×10^{19} eV. This change by a factor of 2 in the energy changes the loss length by almost one order of magnitude, which translates approximately into the same ratio between the flux below and above 5×10^{19} eV if the sources have no luminosity evolution and no local overdensity and there is no magnetic field. A luminosity evolution that makes sources at high redshift brighter than nearby sources enhances this jump, making the feature more pronounced. A local overdensity of sources on the other hand, has the opposite effect. Moreover, hard

spectra contain more particles in the high energy part, therefore they correspond to a GZK feature which is less pronounced than that due to a soft injection spectrum. All these effects were discussed at length for the first time in ¹⁰.

In addition to all these effects, there are some observational issues to take into account: all experiments currently operating have a $\sim 30\%$ statistical error in the energy determination and future experiments will probably not make a big improvement in this respect. Since the spectrum of cosmic rays is steep, there is a larger number of events that are given a higher energy than viceversa, which makes the observed GZK feature look smoother. The effect of a systematic error in the energy determination clearly depends on which way it goes: as shown in ⁴, such an error can have important implications in the comparison between different experiments.

From all this follows that the sharpness of the GZK feature depends on many still unknown parameters. It is likely that the GZK feature is indeed rather smooth, which implies a substantial number of events above 10^{20} eV. This is important for the assessment of the role of next generation UHECR experiments. As an example, in Fig. 2 I plot the expected spectra of UHECRs for the Auger (left panel) and EUSO (right panel) statistics of events, as simulated in ⁴.




Figure 2. Simulated spectra for the Auger (left) and EUSO (right) expected statistics of events as obtained in ².

The injection spectrum used in Fig. 2 is $E^{-2.6}$, which fits the low energy (AGASA and HiRes) data in the case that there is no evolution of the source

luminosity with the redshift. Even a comparison by eye of the AGASA data with these simulated spectra shows that the Auger and EUSO statistics are needed in order to finally settle the issue of the presence of the GZK feature and its detailed shape.

3.2. *What are small scale anisotropies and large scale isotropy telling us?*

Despite the absence of any correlation between the arrival directions of UHECRs and any local matter overdensity (e.g. the Galaxy or the local supercluster), some anisotropies on the scale of a few degrees have been found in the AGASA data ⁵ in the form of doublets and triplets of events. For a truly diffuse distribution of sources these multiplets should appear only as a result of chance coincidences. This possibility appears to be disfavored, although some discussion is still ongoing ⁶ mainly concerning the way the data are treated for the analysis of the statistical significance. In Hires data there seems to be no indication of such anisotropies, which is not surprising for two reasons: 1) for the data obtained in the *mono* regime, the directions of arrival are obtained in the form of ellipses for which the definition of clusters of events is problematic. The data collected in the *stereo* mode on the other hand are fewer; 2) the acceptance of HiRes is energy dependent in the energy region above 10^{19} eV. This means that the number of events needs to be corrected in order to obtain the spectrum of UHECRs, but this correction cannot account for the directions of arrival of the events that the correction is made for.

Clustering of events on small angular scales is the signature that UHECRs are accelerated at astrophysical sources. Moreover, the number of multiplets with different multiplicities is an index of the density of sources, which makes small scale anisotropies a very important tool to have the first important clue to the nature of the sources.

Some attempts to estimate the number of sources of UHECRs in our cosmic neighborhood from the small scale anisotropies found by AGASA have been carried out, adopting both semi-analytical and numerical approaches. An analytical tool to evaluate the chance coincidence probability for arbitrary statistics of events was proposed in ¹¹. A rigorous analysis of the clusters of events and of their energy dependence was given in ¹². In ¹³ the authors use an analytical method to estimate the density of the sources of UHECRs restricting their attention to the 14 events with energy above 10^{20} eV with one doublet. They obtain a rather uncertain estimate centered around $6 \times 10^{-3} \text{ Mpc}^{-3}$. In ¹⁴ the energy losses are introduced through a function, derived numerically, that provides the probability of arrival of a particle from a source at a given distance. Again, only events above 10^{20} eV are considered, therefore the analysis is based upon one doublet of events out of 14 events. This causes extremely large uncertainties in the estimate of the source density, found to be $180_{-165}^{+2730} \times 10^{-3} \text{ Mpc}^{-3}$. No account of the statistical errors in the energy determination nor of the declination dependence of the acceptance

of the experimental apparatus is included in all these investigations. A complete calculation of the number density and luminosity of the sources of UHECRs with a full numerical simulation of the propagation was carried out in ¹⁵. Such a calculation also accounted for the statistical errors in the energy determination and the declination dependence of the acceptance of the experiments involved.




Figure 3. Small scale anisotropies simulated in ¹³ for the AGASA statistics. The thick dots represent the number of doublets and triplets actually observed.

In Fig. 3 I plot the average frequency of occurrence of multiplets with different multiplicities using the AGASA statistics of events, as obtained in ¹⁵. The thick dots represent the number of doublets and triplets observed by AGASA. The panels refer to source density 10^{-6} Mpc^{-3} , 10^{-5} Mpc^{-3} , 10^{-4} Mpc^{-3} from top to bottom, and are obtained for an injection spectrum $E^{-2.6}$ with no redshift evolution (left panels) and injection spectrum $E^{-2.4}$ with redshift evolution $\propto (1+z)^4$ (right panels). The error bars are obtained by simulating many realizations of the source distribution at fixed average source density. The best fit seems to be obtained for a source density $n_s \approx 10^{-5} \text{ Mpc}^{-3}$, although the error bars are large due to the limited statistics of clustered events in AGASA. In ¹⁵ the simulation was extended to mock the statistics expected for Auger and EUSO.

In Fig. 4 I plot the two-point correlation function for the simulated data of Auger (left panel) and EUSO (right panel) ¹⁵. The very pronounced peak at small angular




Figure 4. Two point correlation function for the simulated Auger (left) and EUSO (right) data, as obtained in ¹³ for an angular resolution of 2 degrees. The Auger data refer to energies above 4×10^{19} eV, while the EUSO data refer to energies larger than 10^{20} eV, where the acceptance is expected to be independent of energy.

separation is the evidence for clusters of events, while an approximately flat line would be expected in the case of perfect isotropy down to small scales. It is therefore clear that Auger and EUSO will definitely pin down the density of the sources of UHECRs (The figures were obtained for a source density of 10^{-5} Mpc^{-3}). The wiggles in the EUSO two-point correlation function appear because of statistical fluctuations but their average amplitude is related to the number density of sources ¹⁵. These features start to appear when the exposure of the experiment gets close to the so-called *critical exposure*, $\sigma_c = 42000 (n_s/10^{-5} \text{Mpc}^{-3}) \text{ km}^2 \text{ sr yr}$, defined as the exposure that allows at least one event with energy above 10^{20} eV to reach the detector from each single source within a distance equal to the loss length of these particles.

On large scales there is no evidence for anisotropies in the data. It has been argued that since there is a clear inhomogeneity in the nearby distribution of galaxies (local supercluster), the complete isotropy at energies above 4×10^{19} eV would suggest the presence of a strong local magnetic field ¹⁶. In fact, as discussed at length in ^{4,15}, particles with energy around 4×10^{19} can reach us from distances of 800 – 1000 Mpc, therefore the local inhomogeneous distribution of sources does not reflect into the anisotropy of the arrival directions at this energy (to some extent, this appears to be the conclusion reached later in ¹⁷). In other words, at given

energy, the observed anisotropy would reflect the inhomogeneity of the universe on scales comparable with the loss length at that energy.

3.3. *What is the role of intergalactic magnetic fields on the propagation of UHECRs?*

Our knowledge of the intergalactic magnetic field is very poor and in order to give a fair answer to this question all possibilities should be investigated, provided no observational bound is violated. Magnetic fields of $0.1 - 1 \mu\text{G}$ have been found in virialized structures, such as galaxies and clusters of galaxies. These sources however occupy a fraction $\sim 10^{-5}$ of the volume of the universe and do not play an important role for the propagation of the bulk of UHECRs. The magnetic fields that we are most interested in are therefore the fields in the voids and in non-virialized large scale structures, that occupy most of the volume of the universe. If the magnetic field in the universe is the result of the pollution of astrophysical sources, these regions are expected to be not much magnetized: most field should be where sources are, for the same reason why gas is mostly where sources are. If however the field has a cosmological origin, then even the voids and filamentary regions may have appreciable magnetization. Observationally, the most stringent limits come from Faraday rotation measurements (FRM): these limits depend on assumptions on the topology of the field and on the density of matter along a line of sight. Current limits vary wildly depending on the assumptions adopted, the most severe limit being $B < 10^{-9} - 10^{-10}$ G (e.g. Kronberg¹⁸, but correcting his limit for the fact that Ω_b , the baryon fraction in the universe, is not unity. See¹⁹ for a discussion). Charged particles in a turbulent field perform a random walk. The departure of the trajectory from a straight line stays smaller than some given angular size θ_{exp} if

$$B < 10^{-10} \left(\frac{E}{4 \times 10^{19} \text{eV}} \right) \left(\frac{\theta_{exp}}{2^\circ} \right) \left(\frac{D}{1000 \text{Mpc}} \right)^{-1/2} \left(\frac{L_c}{1 \text{Mpc}} \right)^{-1/2} \text{ Gauss.}$$

Here we can interpret the angle θ_{exp} as the angular resolution of an experiment for the detection of UHECRs. This expression suggests that particles with energy $\sim 4 \times 10^{19}$ eV move in approximate straight line propagation provided the field on the scale specified by the distance D remains smaller than $\sim 10^{-10}$ G. This result is confirmed by recent numerical simulations of the propagation of UHECRs in the magnetized large scale structures of the universe²⁰. For lower energy particles the deflection may become large and the propagation may approach the diffusive regime (see²¹ for a discussion).

The limit imposed by FRM becomes even less stringent on the field in local structures such as the local supercluster. In this case however one can use some physical insights in order to get a hint on the strength of the field: at the present cosmic time, clusters of galaxies are the largest virialized regions in the universe. Larger structures, such as superclusters, did not get to that stage as yet, and are

therefore expected to have smaller velocity dispersion and smaller magnetic fields. The magnetic fields measured in clusters of galaxies (in fact in only some of them) is of order $0.1 - \text{few } \mu\text{G}$, roughly $10 - 100$ times smaller than the equipartition fields ^a (despite the fact that these are virialized structures). In general, if R_{size} is the size of a structure and M the mass in the volume $(4/3)\pi R_{size}^3$, the condition $\frac{B^2}{8\pi} \ll \rho v^2$ translates into

$$B \ll 8\pi \left(\frac{G}{3}\right)^{1/2} R_{size} (\Omega_b \Omega_m)^{1/2} \rho_{cr} \approx 10^{-7} \left(\frac{R_{size}}{10 \text{ Mpc}}\right) \left(\frac{h}{0.7}\right)^{3/2} G,$$

where h is the dimensionless Hubble constant and I used $\Omega_b = 0.02h^{-2}$ for the baryon fraction and $\Omega_m = 0.3$ for the mass fraction.

3.4. *Where and how are UHECRs accelerated?*

A first piece of the answer to this question may come from the measurement of the spectral shape in the energy region where we expect to find the GZK feature and from the measurement of the chemical composition. For instance a weak or absent GZK feature, together with an appreciable fraction of gamma rays in the composition would point toward a top-down origin of UHECRs. In this case it is difficult to envision how prominent small scale anisotropies may arise. In the following, due to the limited space available I will not discuss these models any further (see ²² for a review of this class of models).

As discussed above, both the shape of the GZK feature and the presence of small scale anisotropies should tell us about the source density and luminosity. For sources which are continuous in time, as stressed above, the source density inferred from SSA is $\sim 10^{-5} \text{ Mpc}^{-3}$, corresponding to a source luminosity $\sim 10^{42} \text{ erg s}^{-1}$ at energies above 10^{19} eV .

A discussion of 'Where' may UHECRs be accelerated can be found in many excellent reviews ^{23,24,25,26} and the limited space available here does not allow me to add much to what is discussed there. In the following I will therefore concentrate on two issues concerning the 'How' UHECRs are accelerated. More specifically I will spend a few words on two subjects, namely non-linear shock acceleration at newtonian shocks, and particle acceleration at relativistic shocks that are particularly important for the acceleration of UHECRs.

Most scenarios that have been proposed so far for the acceleration of UHECRs in astrophysical sources, with few exceptions, are related to particle acceleration at either newtonian or relativistic shock waves. It is therefore important that we understand the details of this process in order to confront observations. A typical assumption adopted to estimate the plausibility of a class of sources is that the spectrum of accelerated particles is a power law and the energy it contains is a

^aOne should keep in mind that the concept of equipartition field here is different from the one typically used in radio astronomy, in which the equipartition is meant with the radiating relativistic particles. The latter field is clearly quite smaller than the former.

small fraction of the total energy budget. It has been found in recent times that neither one of these assumptions need to be valid. Numerical calculations suggest²⁷ that although the fraction of particles accelerated to suprathermal energies is in fact small, the energy channelled into these few particles may be a substantial part of the kinetic energy flux through the shock surface. This effect is due to the backreaction of the accelerated particles on the shock itself, that enhances the efficiency of acceleration instead of reducing it. When this happens, the spectrum of the accelerated particles is no longer a power law, as shown also in analytical calculations of the nonlinear backreaction^{28,29}. More specifically, for strongly modified shocks the spectra are such that most energy is concentrated at p_{max} , the maximum momentum achievable at the shock, something which is at odds with the predictions of the linear theory, where the energetics of accelerated particles is typically dominated by the mildly relativistic particles. It is easy to envision the consequences that these findings may have for the acceleration of UHECRs, since in this case there is the constant need to channel as much energy as possible in the highest energy part of the spectrum of a source.

In some candidate sources of UHECRs, such as gamma ray bursts, the acceleration is postulated to occur at relativistic shock waves. The spectrum of accelerated particles in these cases depends quite sensibly on the scattering properties of the media upstream and downstream of the shock. In most cases however, the spectra of accelerated particles are calculated in the assumption of small pitch angle scattering and are found to be power laws with a slope that for ultra-relativistic shocks tends to be almost universal³⁰, $\gamma \sim 2.2 - 2.4$. Recently in³¹ a new approach has been proposed that describes shock acceleration at shocks with arbitrary velocity (from newtonian to relativistic) and arbitrary scattering properties of the medium. In³², the approach was further developed and checked versus several test cases in both the newtonian and relativistic regimes. The approach allows one to determine analytically the spectrum of particles and the anisotropy in pitch angle even for those situations in which the assumption of small pitch angle scattering does not apply.

Some general comments on shock acceleration may be made independently of the details: the acceleration of particles at ultra-relativistic shocks is likely to proceed in two steps. During the first step, a factor $\sim \Gamma^2$ can be gained by the particles (Γ here is the Lorentz factor of the shock). After this first step, the distribution of particles upstream is beamed within a cone of aperture $1/\Gamma$ with respect to the direction of motion of the shock and the energy gain is only of order unity in the successive shock crossings. The acceleration at relativistic shocks requires some appreciable level of turbulence in the downstream fluid: in such frame, the shock moves with a speed $\sim 1/3$, therefore the particles would not be able to reach the shock if the field is coherent on large scales. A recent review of shock acceleration at relativistic shocks can be found in³³.

4. Conclusions

I discussed some general issues related to the origin and propagation of UHECRs, trying to point out which future developments may contribute to improve our understanding of the current problems.

References

1. K. Greisen, Phys. Rev. Lett. 16 (1966) 748; G. T. Zatsepin and V. A. Kuzmin, Sov. Phys. JETP Lett. 4, 78 (1966).
2. M. Takeda et al., Phys. Rev. Lett. 81, 1163 (1998); M. Takeda et al., preprint astro-ph/9902239; N. Hayashida et al., Phys. Rev. Lett. 73, 3491 (1994); D.J. Bird et al. Astrophys J. 441, 144 (1995); Phys. Rev. Lett. 71, 3401 (1993); Astrophys. J. 424, 491 (1994); M.A. Lawrence, R.J.O. Reid and A.A. Watson, J. Phys. G. Nucl. Part. Phys. 17, 773 (1991) 773; N.N. Efimov et al., Ref. Proc. International Symposium on *Astrophysical Aspects of the Most Energetic Cosmic Rays*, eds. M. Nagano and F. Takahara (World Scientific, Singapore, 1991), p. 20
3. T. Abu-Zayyad, et al., preprint astro-ph/0208301; T. Abu-Zayyad, et al., preprint astro-ph/0208243.
4. D. De Marco, P. Blasi, and A.V. Olinto, Astropart. Phys. **20**, 53 (2003).
5. M. Takeda, et al., Astrophys. J. **522**, 225 (1999).
6. C.B. Finley, and S. Westerhoff, preprint astro-ph/0309159.
7. M. Ave, J.A. Hinton, R.A. Vazquez, A.A. Watson, A.A., and E. Zas, Phys. Rev. **D65**, 063007 (2002).
8. A.A. Watson, preprint astro-ph/0312475 (these proceedings).
9. V. Berezhinsky, A. Z. Gazizov and S. Grigorieva, preprint astro-ph/0210095; V. Berezhinsky, A. Z. Gazizov and S. Grigorieva, preprint hep-ph/0204357.
10. V. Berezhinsky and S. Grigorieva, Astron. Astroph. **199**, 1 (1988).
11. H. Goldberg and T.J. Weiler, Phys. rev. **D64**, 056008 (2001).
12. W.S. Burgett and M.R. O'Malley, Phys. Rev. **D67**, 092002 (2003).
13. S.L. Dubovsky, P.G. Tinyakov and I.I. Tkachev, Phys. Rev. Lett. **85**, 1154 (2000).
14. Z. Fodor and S.D. Katz, Phys.Rev. **D63**, 023002 (2001).
15. P. Blasi, and D. De Marco, Astropart. Phys., in press (preprint astro-ph/0307067).
16. G. Sigl, F. Miniati, and T. Ensslin, Phys.Rev. **D68**, 043002 (2003).
17. G. Sigl, F. Miniati, and T. Ensslin, preprint astro-ph/0309695.
18. P.P. Kronberg, Reports on Progress in Physics **57**, 325 (1994).
19. P. Blasi, S. Burles and A.V. Olinto, Astrophys. J. Lett. **514**, 79 (1999).
20. K. Dolag, D. Grasso, V. Springel and I. Tkachev, preprint astro-ph/0310902.
21. T. Stanev, preprint astro-ph/0303123.
22. P. Bhattacharjee and G. Sigl, Phys. Rept. **327**, 109 (2000).
23. A.V. Olinto, Phys. Rep. **333**, 329 (2000).
24. F.W. Stecker, J. Phys. **G29**, 47 (2003).
25. R.J. Protheroe and R.W. Clay, preprint astro-ph/0311466.
26. P. Blasi, Proceedings of the XXI Texas Symposium on Relativistic Astrophysics (Editors: R. Bandiera, R. Maiolino and F. Mannucci), p. 387 (2003).
27. H. Kang, T.W. Jones and U.D.J. Gieseler, Astroph. J. **579**, 337 (2002).
28. M.A. Malkov, Astrop. J. **491**, 584 (1997); M.A. Malkov, P.H. Diamond, and H.J. Völk, Astroph. J. **533**, 171 (2000).
29. P. Blasi, Astropart. Phys. **16**, 429 (2002); P. Blasi, Astropart. Phys. in press (preprint astro-ph/0310507).

30. A. Achterberg, Y.A. Gallant, J.G. Kirk, and A.W. Guthmann, MNRAS **328**, 393 (2001).
31. M. Vietri, *Astroph. J.* **591**, 954 (2003).
32. P. Blasi and M. Vietri, *On particle acceleration around shocks II: a fully general method for arbitrary shock velocities and scattering media*, submitted to *Astroph. J.*
33. M. Ostrowski, *Journal of Physical Studies* **6**, 393 (2002).